

Bulk viscous effects near the QCD critical point

Yi Yin





A. Monnai, S. Mukherjee and YY, in preparation

Opportunities for Exploring Longitudinal Dynamics in Heavy Ion Collisions


RBRC workshop, BNL, Jan. 20th, 2016

Highlights of the talk

- Bulk viscous pressure is very sensitive to the presence of the QCD critical point. $\Pi \sim \xi^3$
- Sizable critical fluctuations should be accompanied with influence due to enhanced bulk viscous pressure.
- Exploratory study with I+I hydro.



QCD phase diagram and the critical point





A sketch of the QCD phase diagram.

- QCD critical point: a distinctive feature in QCD phase diagram. (Its location is *unknown* theoretically).
- Beam Energy Scan Program: search for QCD critical point experimentally.

Correlation length and observables

- Correlation length ξ_{eq} of critical modes universally grows and becomes divergent near a critical point.
- Quantities with strong dependence on correlation length: important for search for QCD critical point.
- Well-studied examples: fluctuation observables such as cumulants of net-proton numbers.
- Cumulants data is consistent with expectation due to critical fluctuation. Any other observables sensitive to the critical point?



Universal critical dynamics of the QCD critical point: I

- Both static properties (fluctuation) and dynamical properties (transport) universally scale with the correlation length.
- Two types of degrees of freedom:
 - short-range: rapid equilibration.
 - long-range (critical mode) : equilibrate diffusively and slowly (critical slowing down).
- Relaxation time of the critical mode: $\tau_\sigma \sim \xi_{eq}^z$ with $z > 0$ dynamic critical exponent.

Universal critical dynamics of the QCD critical point: II

- Dynamical universality class (Hohenberg-Halperin, Rev. Mod. Phys, 1977) : long distance dynamics should be matched to hydrodynamics.
- Universality class of QCD critical point: model H (Son-Stephanov, 2004), i.e. the same dynamic universality as Liquid-Gas transition.

$$z = 3 + \mathcal{O}(\epsilon)$$

- Transport coefficients scale with correlation length:

$$\text{(Baryon) conductivity } \sigma \sim \xi_{\text{eq}},$$

$$\text{diffusive constant } D \sim \xi_{\text{eq}}^{-1},$$

$$\text{shear viscosity } \eta \sim \xi_{\text{eq}}^{\mathcal{O}(\epsilon)}$$

- Bulk viscosity strongly depends on ξ_{eq} : $\zeta \sim \xi_{\text{eq}}^3$ (Onuki, 1997; Karsch-Kharzeev-Tuchi, PLB 2008; Moore-Saremi, JHEP 2008)

This talk: effects of bulk viscous pressure on search for QCD critical point.

- Introduction and motivations.
- **Bulk viscous pressure near the QCD critical point**
- Longitudinal expansion near the QCD critical point
- Summary and outlook

Bulk viscous pressure and bulk viscosity

- Bulk viscous pressure: non-equilibrium contribution to the effective pressure :

$$T^{\mu\nu} = \epsilon u^\mu u^\nu + (p - \Pi) \Delta^{\mu\nu} + \text{shear viscous term}$$

$$p_{eff} \equiv p_{eq} - \Pi$$

- First order hydro. (Navier-Stokes limit):

$$\Pi \rightarrow \Pi_\zeta \equiv \zeta \partial_\mu u^\mu$$

$$\Pi \sim \zeta \stackrel{?}{\sim} \xi_{eq}^3$$

- Growth of bulk pressure is limited by finite time effects: (c.f. Song-Heinz, 2009)

$$(u^\mu \partial_\mu) \Pi \sim \partial_\tau \Pi = \frac{1}{\tau_\Pi} [\Pi_\zeta - \Pi] \quad \text{Israel-Stewart Theory}$$

- Key input: the behavior of τ_Π near a critical point (unknown).

Bulk viscosity and τ_Π near the QCD critical point


$\zeta \sim$ equilibration of p under compression (decompression)

- Pressure depends on the critical mode:
equilibration time of pressure $\sim \tau_\sigma$

$$\zeta \sim \tau_\sigma \sim \xi_{eq}^3$$

$\tau_\Pi \sim$ equilibration of p under a perturbation δp

$$\text{IS Theory} \rightarrow \partial_\tau \delta p = -\frac{\delta p}{\tau_\Pi}$$



- **New** result (A. Monnai, S. Mukherjee and YY, in preparation):


$$\tau_\Pi \sim \tau_\sigma \sim \xi_{eq}^3$$

- Causality constrain (Romatschke. 2009):

$$\frac{dc_s(k)}{dk} \leq 1 \rightarrow \frac{\zeta}{\tau_\Pi} = \text{bounded}$$

A toy model

- Solving relaxation equation along a trajectory passing the critical regime:





$$u^\mu \partial_\mu \Pi = \frac{1}{\tau_\Pi} [\zeta (\partial_\mu u^\mu) - \Pi] \rightarrow \partial_\tau \Pi = \frac{1}{\tau_\Pi} \left[\zeta \left(\frac{n_V}{\tau} \right) - \Pi \right]$$

- Implementing critical dynamics:

$$\zeta = \zeta_0 \left(\frac{\xi_{\text{eq}}}{\xi_0} \right)^3$$

$$\tau_\Pi = \tau_{\Pi,0} \left(\frac{\xi_{\text{eq}}}{\xi_0} \right)^3$$

Evolution of bulk viscous pressure



Evolution of non-equilibrium correlation length (Berdnikov-Rajagopal, 2000)

$$\partial_\tau \xi = (\xi_{\text{eq}} - \xi) / \tau_\sigma$$

- Finite time effects:
 - delay the growth of Π .
 - Preserve critical behavior.
- Similar to the evolution of non-equilibrium length.

Bulk viscous pressure and (non-equilibrium) correlation length



- The growth of ξ is accompanied with the growth of Π (tested numerically) .

$$\Pi \sim \xi^3$$

- The critical behavior $\zeta \sim \tau_{\Pi} \sim \tau_{\sigma} \sim \xi_{\text{eq}}^3$ are controlled by the same physics. (critical slowing down).

- Important implications:

Sizable critical fluctuations in data



Sizable effects due to Π

Observables sensitive to bulk viscous pressure

- Enhanced bulk pressure will reduce the effective pressure :

$$p_{eff} \equiv p_{eq} - \Pi$$

- If Π is too large thus $p_{eff} < 0$: cavitation (hydro. is broken)
- Within hydro. (IS theory): influence flow.

In fluid rest-frame

$$\partial_\mu T^{\mu\nu} = 0$$



$$\partial_t \vec{v} \propto \nabla p_{eff}$$

- Particle distribution at freeze-out:

$$\delta f \propto \frac{p_T^2}{T^2} \frac{\Pi}{\epsilon + p}$$

- Quantitative study: hydro. simulation with critical dynamics.


- Introduction and motivations.
- Bulk viscous pressure near the QCD critical point
- Longitudinal expansion near the QCD critical point
- Summary and outlook

Set-up

- First attempt to study bulk viscous effects near the QCD critical point (A. Monnai, S. Mukherjee and YY, in preparation).
- I + I Israel-Stewart theory including baryon density.
 - E.O.S: lattice QCD with Taylor expansion.
 - Initial condition: CGC (energy density) + Valence quark dist. (baryon density).
 - Linear mapping to Ising model

Preliminary Results I

- Run at 17 GeV, *assuming* (cross-over) side of the critical regime is probed. (Disclaimer: neither fitting the data or making prediction).
- Three different inputs:
 - Ideal hydro.
 - Bulk term without critical enhancement
 - Bulk term + critical enhancement



$$\zeta = \zeta_0 \left(\frac{\xi_{\text{eq}}}{\xi_0} \right)^3$$


$$\tau_{\Pi} = \tau_{\Pi,0} \left(\frac{\xi_{\text{eq}}}{\xi_0} \right)^3$$

Bulk viscous pressure and longitudinal flow

$$\eta = 1$$

- Enhanced bulk viscous pressure in the critical regime.
- Change of flow is frozen.

$$\partial_t \vec{v} \propto \nabla p_{eff}$$
- Particle distribution, in progress.



- Introduction and motivations.
- Bulk viscous pressure near the QCD critical point
- Longitudinal expansion near the QCD critical point
- **Summary and outlook**

Summary

- Explore the critical dynamics and its implication on search for QCD critical point.
- Bulk viscous pressure is correlated and sensitive to the growth of correlation length.
- Flow observables: sensitive to the reduction of effective pressure. (Complementary to fluctuation observables, higher statistics)
- Exploratory study with I+I hydro underway.

Outlook

- Directed flow observables: reaches minimum for smaller pressure (the softest point due to bulk viscous pressure?).
- Improve the domain of validity of hydro near a QCD critical point (in progress with M. Stephanov).
- Critical behavior of second order hydro. coefficients (extension of the classic work by Hohenberg-Halperin).
- Many interesting questions ahead!

$$\partial_t \vec{v} \propto \nabla p_{eff}$$

